

Dax Varkey MD, MPH Caroline McBane PA-C 1130 N. Church Street, Suite 100 336-375-2300 (tel) 336-375-2314 (fax)

Patient:	Date of Surgery:		
Procedure: Right/Left Rotator Cuff Repair Associated procedures: Biceps Tenodesis: avoid resisted elbow flexion x 6 weeks Distal Clavicle Resection: Avoid cross-body adduction x 6 weeks First week: Sling at all times except for hygiene and pendulums if instructed. Home exercises of elbow and wrist ROM, grip strengthening and small pendulums.			
		Weight bea Brace: Sling Ice: Not dire for first 2 we ROM: PROM max.	Begin formal PT (2-3x per week) ring: NWB, ok for typing if in sling at all times except hygiene. Pillow is optional for sling. ctly on skin, recommend as much as possible. 5x/day, 20 min/session eks. Then after activity at minimum. If only – goal of forward flexion 140°, ER at side 40°, abduction 60° deg
		Weight bea Brace: Disc ROM: Light and progres 12 weeks. Exercises: level. Deltoi	cing: < 2-5 lbs operative arm ontinue sling after 6 weeks. Deassive stretching at end ranges. Begin AAROM (canes, pulleys, etc.) is supine to vertical. Gradually progress with AROM. Goal full motion by Begin periscapular, pec/lat/trapezius isometrics with arm below shoulder d and cuff isometrics with arm at side at 8 weeks. Begin light g at 10-12 weeks at therapist discretion.
Weight bea ROM: No re Exercises: A reps x 2-3 se	B wks): Progress with strength ring: Advance to tolerance. Striction Advance as tolerated to bands and to light weights (1-5 lbs) with 8-12 ets for cuff, deltoid, scapular stabilizers. No more than 3x / week to avoid so. Begin to eccentrically resisted motions.		
Weight bea Sports: At 5 Work: Overl take 4-6 mo	mo ok to progress to light throwing. Golf may begin at 4 months. nead heavy lifting can begin to tolerance at ~4 months. Heavy labor may		
Frequency:	x/week xweeks		
By signing this referral, I certify that I have examined this patient and physical therapy is medically necessary.			
-	Date:		

v 22 10 28

Dax Varkey MD, MPH